

ON p-ADIC FUNCTIONAL INTEGRATION

Goran S. DJORDJEVIĆ, Branko DRAGOVICH*

*Department of Physics, University of Niš,
P.O.Box 91, 18001 Niš, Yugoslavia
E-mail: gorandj@junis.ni.ac.yu*

**Institute of Physics, P.O.Box 57, 11001
Belgrade, Yugoslavia
E-mail: dragovic@phy.bg.ac.yu*

Abstract

p-Adic generalization of the Feynman path integrals in quantum mechanics is considered. The probability amplitude $\mathcal{K}_v(x'', t''; x', t')$ ($v = \infty, 2, 3, \dots, p, \dots$) for a particle in a constant field is calculated. Path integrals over $\mathbb{Q}_p$ have the same form as those over $\mathbb{R}$.

1. Introduction

This investigation of p-adic functional integration is motivated by a successful application of p-adic numbers in many parts of theoretical and mathematical physics, and particularly in quantum mechanics (for a review see Refs. 1,2,3). As it is known [4], p-adic numbers may be obtained by completion of the field of rational numbers $\mathbb{Q}$ with respect to the p-adic norm (valuation). Recall that the field of real numbers $\mathbb{R}$ may be regarded as completion of $\mathbb{Q}$ with respect to the usual absolute value.

In physical measurements only rational numbers can be obtained, and comparison of theoretical models and experimental results performs within $\mathbb{Q}$. Since there is a good analysis on $\mathbb{R}$ theoretical models are traditionally constructed over $\mathbb{R}$ and not over $\mathbb{Q}$. Only in 1987 the use of p-adic analysis started to give acceptable physical models.

The fields of p-adic numbers $\mathbb{Q}_p$ (p denotes any prime number) and $\mathbb{R}$ exhaust all possible number fields which can be obtained by completion of $\mathbb{Q}$. It is a natural hope that inclusion of p-adic numbers in the study of physical phenomena provides some additional clarification of a problem in question. The space of adeles is a mathematical concept which

contains all $\mathbb{Q}_p$ and $\mathbb{R}$, and enables to treat them simultaneously.

One of the greatest achievements in the use of p-adic numbers in physics is a formulation of p-adic quantum mechanics [5]. The elements of the corresponding Hilbert space $L_2(\mathbb{Q}_p)$ are complex-valued functions of a p-adic argument. It is recently shown [6] that p-adic quantum mechanics and the ordinary one can be unified in the form of adelic quantum mechanics, which is quantum mechanics over the space of adeles.

One of the basic characteristics of quantum mechanics is a probabilistic description of physical events. As a starting point to calculate quantum dynamics, instead of the Schrödinger equation, one can use the kernel $\mathcal{K}(x'', t''; x', t')$ of the evolution operator $U(t)$. This $\mathcal{K}(x'', t''; x', t')$ is also called the quantum-mechanical propagator, or the probability amplitude for a particle to go from a space-time point (x', t') to a space-time point (x'', t'') . For the reason of simplicity we consider only 1+1 - dimensional systems.

Feynman has postulated [7] $\mathcal{K}(x'', t''; x', t')$ to be the path integral

$$\mathcal{K}(x'', t''; x', t') = \int \exp\left(\frac{2\pi i}{h} S[q]\right) \mathcal{D}q, \quad (1.1)$$

where $S[q] = \int_{t'}^{t''} L(q, \dot{q}, t) dt$ is an action. $L(q, \dot{q}, t)$ is the Lagrange function and $x'' = q(t'')$, $x' = q(t')$. The integral in (1.1) is a symbol which has to realize an intuitive understanding that a quantum-mechanical particle can propagate from x' to x'' using uncountably many paths which connect these two points. Thus the Feynman path integral means a continual summation of single amplitudes $\exp(\frac{2\pi i}{h} S[q])$ over all paths $q(t)$ staying between $x' = q(t')$ and $x'' = q(t'')$. In practical calculations it is the limit of an ordinary multiple integral over $N - 1$ variables $q_i = q(t_i)$ when $N \rightarrow \infty$. Namely, the time interval $t'' - t'$ is divided into N equal subintervals and integration is taken over q_i from $-\infty$ to $+\infty$, but for a fixed time t_i .

The kernel $\mathcal{K}$ defined in (1.1) must satisfy the following conditions:

$$(i) \quad \mathcal{K}(x'', t''; x', t') = \int_R \mathcal{K}(x'', t''; x, t) \mathcal{K}(x, t; x', t') dx \quad (1.2)$$

$$(ii) \quad \int_R \mathcal{K}^*(x'', t''; x, t) \mathcal{K}(x', t''; x, t) dx = \delta(x'' - x') \quad (1.3)$$

$$(iii) \quad \mathcal{K}(x'', t; x', t) = \delta(x'' - x'). \quad (1.4)$$

For the classical action $\bar{S}(x'', t''; x', t')$, *i.e.* the minimal action with the classical trajectory $\bar{q} = \bar{q}(t)$, which is quadratic in x'' and x' , it has been shown [8] that in the real

case

$$\mathcal{K}(x'', t''; x', t') = \left(\frac{i}{h} \frac{\partial^2 \bar{S}}{\partial x'' \partial x'} \right)^{\frac{1}{2}} \exp \left(\frac{2\pi i}{h} \bar{S}(x'', t''; x', t') \right), \quad (1.5)$$

where h is the Planck constant.

Feynman invented path integral in 1942, but he published it in 1948 [7]. Since then it has been a subject of permanent interest in theoretical and mathematical physics, and it is now one of the best approaches to quantum theory. On the recent progress, present state and some future prospects one can see Ref. 9.

This paper is devoted to the p-adic generalization of (1.1), *i.e.* $x'', x'; t'', t'$ are p-adic variables and $\mathcal{K}$ is the complex-valued function. Such approach is suggested in Ref. 5 and its validity is demonstrated on the harmonic oscillator [10]. We shall see here that it can be formally done in the same way, and that it has the same form, as in the real case.

2. On p-adic numbers and p-adic analysis

We recall here only some basic properties which are relevant to the rest of the paper.

Any $x \in \mathbb{Q}_p$ can be presented in the form

$$x = p^\nu (x_0 + x_1 p + x_2 p^2 + \dots), \quad \nu \in \mathbb{Z}, \quad (2.1)$$

where $x_i = 0, 1, \dots, p-1$ are digits. p-Adic norm of any term $x_i p^{\nu+i}$ in the canonical expansion (2.1) is $|x_i p^{\nu+i}|_p = p^{-(\nu+i)}$. Since p-adic norm is the non-archimedean one, *i.e.* $|a+b|_p \leq \max\{|a|_p, |b|_p\}$, it follows that $|x|_p = p^{-\nu}$. p-Adic numbers are closely connected to the prime numbers ($p = 2, 3, 5, \dots$) and there are infinitely many $\mathbb{Q}_p$, *i.e.* for every p there is one corresponding $\mathbb{Q}_p$.

There is no natural ordering on $\mathbb{Q}_p$. However one can introduce a linear order on $\mathbb{Q}_p$ by the following definition: $x < y$ if $|x|_p < |y|_p$ or when $|x|_p = |y|_p$ there exists such index $m \geq 0$ that digits satisfy $x_0 = y_0, x_1 = y_1, \dots, x_{m-1} = y_{m-1}, x_m < y_m$.

There are two kinds of analysis over $\mathbb{Q}_p$: the first one which is connected with map $\varphi : \mathbb{Q}_p \rightarrow \mathbb{Q}_p$, and another one related to map $f : \mathbb{Q}_p \rightarrow \mathbb{C}$, where $\mathbb{C}$ is the field of complex numbers. Here, we use the both of these analysis. Derivatives of $\varphi(x)$ are defined as in the real case, but with respect to the p-adic norm. There is no integral $\int \varphi(x) dx$ in a sense of the Lebesgue measure [4], but one can introduce $\int_a^b \varphi(x) dx = \Phi(b) - \Phi(a)$ as a functional of analytic functions $\varphi(x)$, where $\Phi(x)$ is an antiderivative of $\varphi(x)$. In the case of map $f : \mathbb{Q}_p \rightarrow \mathbb{C}$ there is well-defined Haar integral. We use here the Gauss integral

$$\int_{\mathbb{Q}_p} \mathcal{X}_v(ax^2 + bx) dx = \lambda_v(a) |2a|_v^{-\frac{1}{2}} \mathcal{X}_v\left(-\frac{b^2}{4a}\right), \quad a \neq 0, \quad (2.2)$$

where index v denotes real and any of p-adic cases, *i.e.* $v = \infty, 2, 3, 5, \dots$, where ∞ corresponds to the real case. $\mathcal{X}_v$ is a multiplicative character: $\mathcal{X}_\infty(x) = \exp(-2\pi i x)$, $\mathcal{X}_p(x) = \exp(2\pi i \{x\}_p)$ where $\{x\}_p$ is the fractional part of $x \in \mathbb{Q}_p$. Function $\lambda_v(a)$ is the complex-valued one defined as follows [1]:

$$\lambda_\infty(a) = \frac{1}{\sqrt{2}}(1 - i \operatorname{sign} a) , \quad (2.3)$$

$$\lambda_p(a) = \begin{cases} 1, & \nu = 2k, \quad p \neq 2, \\ \left(\frac{a_0}{p}\right), & \nu = 2k+1, \quad p \equiv 1 \pmod{4}, \\ i\left(\frac{a_0}{p}\right), & \nu = 2k+1, \quad p \equiv 3 \pmod{4}, \end{cases} \quad (2.4)$$

$$\lambda_2(a) = \begin{cases} \frac{1}{\sqrt{2}}[1 + (-1)^{a_1}i], & \nu = 2k, \\ \frac{1}{\sqrt{2}}(-1)^{a_1+a_2}[1 + (-1)^{a_1}i], & \nu = 2k+1, \end{cases} \quad (2.5)$$

where p-adic number a has the form (2.1), $\left(\frac{a_0}{p}\right)$ is the Legendre symbol, and $k \in \mathbb{Z}$. We will mainly use the following properties of $\lambda_v(a)$:

$$\begin{aligned} \lambda_v(0) &= 1, \quad \lambda_v(a^2b) = \lambda_v(b), \quad \lambda_v(a)\lambda_v(b) = \lambda_v(a+b)\lambda_v\left(\frac{1}{a} + \frac{1}{b}\right), \\ \lambda_v^*(a)\lambda_v(a) &= 1. \end{aligned} \quad (2.6)$$

3. p-Adic generalization of the Feynman path integral

Let us now consider

$$\mathcal{K}_v(x'', t''; x', t') = \int \mathcal{X}_v\left(-\frac{1}{h}S[q]\right) \mathcal{D}_v q \quad (3.1)$$

as a generalization of (1.1) in ordinary quantum mechanics on all number fields $\mathbb{Q}_v$ which can be obtained by completion of $\mathbb{Q}$ (note that $\mathbb{Q}_\infty \equiv \mathbb{R}$). In (3.1) $\mathcal{K}_v$ is a complex-valued function of $x'', t'', x', t' \in \mathbb{Q}_v$ ($v = \infty, 2, 3, \dots$).

DEFINITION. If there exists the limit

$$\begin{aligned} \lim_{N \rightarrow \infty} \mathcal{K}_v^{(N)}(x'', t''; x', t') &= \lim_{N \rightarrow \infty} A_N \int_{\mathbb{Q}_v} \cdots \int_{\mathbb{Q}_v} \\ \mathcal{X}_v\left(-\frac{1}{h} \sum_{i=1}^N \bar{S}(q_i, t_i; q_{i-1}, t_{i-1})\right) & dq_1 \cdots dq_{N-1}, \end{aligned} \quad (3.2)$$

where $x' = q_0$, $x'' = q_N$, $t' = t_0$, $t'' = t_N$, and $A_N = A_N(t'', t')$ is a normalization factor, we say

$$\mathcal{K}_v(x'', t''; x', t') = \lim_{N \rightarrow \infty} \mathcal{K}_v^{(N)}(x'', t''; x', t') \quad (3.3)$$

is the Feynman path integral.

It is understood that $t_0 < t_1 < \dots < t_{N-1} < t_N$ according to linear order on $\mathbb{Q}_v$ and that $|t_i - t_{i-1}|_v \rightarrow 0$ for every $i = 1, 2, \dots, N$ when $N \rightarrow \infty$. Note that $\bar{S}$ in (3.2) is the classical action which corresponds to the classical path $\bar{q}(t)$. This path $\bar{q}(t)$ satisfies the Euler-Lagrange equation

$$\frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} = 0, \quad L = L(q, \dot{q}, t) \quad (3.4)$$

and yields the minimum (extreme) of the action $S[q]$.

4. Generalized path integral for some physical systems

To present time the path integral (3.1) has been derived in the real case for many physical systems [9]. However, p-adic path integral is calculated only for the harmonic oscillator [10]. Here we give the result of calculation for a particle in a constant external field. For the real case this integral can be found in Ref. 7. For the reason of simplicity we use units in which $m = h = 1$, where m is the mass of a particle and h is the Planck constant.

The Lagrange function of a particle with a constant acceleration a is

$$L(q, \dot{q}) = \frac{\dot{q}^2}{2} + aq, \quad (4.1)$$

where q denotes position. The corresponding Euler-Lagrange equation (3.4) is $\ddot{q} = a$. If we denote $t' = 0$, $t'' = T$, $q(0) = q'$, $q(T) = q''$ one obtains the classical path

$$q(t) = \frac{a}{2}t^2 + \frac{q'' - q' - \frac{a}{2}T^2}{T}t + q'. \quad (4.2)$$

For (4.2) one can calculate the classical action

$$\bar{S}(q'', T; q', 0) = \int_0^T \left(\frac{\dot{q}^2}{2} + aq \right) dt \quad (4.3)$$

and it yields

$$\bar{S}(q'', T; q', 0) = \frac{(q'' - q')^2}{2T} + \frac{a}{2}(q'' + q')T - \frac{a^2}{24}T^3. \quad (4.4)$$

Note that expressions (4.1) to (4.4) are valid for real and all p-adic cases.

Let us decompose the time interval T into N subintervals $0 = t_0 < t_1 < \dots < t_{N-1} < t_N = T$ and denote $\varepsilon_i = t_i - t_{i-1}$, $q_i = q(t_i)$ for $i = 1, \dots, N$. To use definition (3.2) let us prove the following

PROPOSITION. It holds

$$\begin{aligned} & \prod_{i=1}^N \lambda_v(2\varepsilon_i) \mid \varepsilon_i \mid v^{-\frac{1}{2}} \int_{\mathbb{Q}_v} \dots \int_{\mathbb{Q}_v} \mathcal{X}_v \left[-\frac{1}{2} \sum_{i=1}^N \frac{(q_i - q_{i-1})^2}{\varepsilon_i} - \frac{a}{2} \sum_{i=1}^N \right. \\ & \left. (q_i + q_{i-1})\varepsilon_i + \frac{a^2}{24} \sum_{i=1}^N \varepsilon_i^3 \right] dq_1 \dots dq_{N-1} = \lambda_v \left(2 \sum_{i=1}^N \varepsilon_i \right) \mid \sum_{i=1}^N \varepsilon_i \mid v^{-\frac{1}{2}} \\ & \mathcal{X}_v \left[-\frac{1}{2} \frac{(q_N - q_0)^2}{\sum_{i=1}^N \varepsilon_i} - \frac{a}{2} (q_N + q_0) \sum_{i=1}^N \varepsilon_i + \frac{a^2}{24} \left(\sum_{i=1}^N \varepsilon_i \right)^3 \right]. \end{aligned} \quad (4.5)$$

In order to prove (4.5) we use the method of induction, integral (2.2) and properties (2.6) of λ_v functions. First we show validity of (4.5) for $N = 2$, and then we assume it is valid for N and show that it holds for $N + 1$.

Denoting $A_N = \prod_{i=1}^N \lambda_v(2\varepsilon_i) \mid \varepsilon_i \mid v^{-\frac{1}{2}}$ we have by (3.2) that $\mathcal{K}_v^{(N)}(x'', t''; x', t')$ for a particle of a constant acceleration is equal to the left-hand side of (4.5). According to the above proposition $\lim_{N \rightarrow \infty} \mathcal{K}_v^{(N)}(x'', t''; x', t')$ is the limit of the right-hand side of (4.5) and it exists. Recall that $q_N = q''$ for any N and $q_0 = q'$, as well as $t'' = T$ and $t' = 0$. Thus we can write down

$$\begin{aligned} & \mathcal{K}_v(x'', T; x', 0) = \lambda_v(2T) \mid T \mid v^{-\frac{1}{2}} \\ & \mathcal{X}_v \left[-\frac{1}{2} \frac{(q'' - q')^2}{T} - \frac{a}{2} (q'' + q')T + \frac{a^2}{24} T^3 \right]. \end{aligned} \quad (4.6)$$

Looking at (4.6) one can see that it has the form

$$\mathcal{K}_v(x'', T; x', 0) = A_v(T) \mathcal{X}_v[-\bar{S}(x'', T; x', 0)]. \quad (4.7)$$

In the real case $A_\infty(T)$ is usually written as $(iT)^{-\frac{1}{2}}$ [7], but one can easily show that it is equal to $\lambda_\infty(2T) \mid T \mid_\infty^{-\frac{1}{2}}$, where $\mid \mid_\infty$ denotes the standard absolute value.

To obtain the corresponding result for a free particle it is enough to put acceleration $a = 0$ in (4.6), *i.e.* for a free particle we get

$$\mathcal{K}_v(x'', T; x', 0) = \lambda_v(2T) \mid T \mid v^{-\frac{1}{2}} \mathcal{X}_v \left[-\frac{1}{2} \frac{(q'' - q')^2}{T} \right], \quad (4.8)$$

and it is in agreement with the known result in the real case [7].

Using the above procedure we also calculated the Feynman path integral for the following physical models:

(i) the de Sitter minisuperspace model [11]

$$\begin{aligned} \mathcal{K}_v(x'', T; x', 0) &= \lambda_v(-2T) \mid 4T \mid_v^{-\frac{1}{2}} \\ \mathcal{X}_v \left[\frac{(q'' - q')^2}{8T} + \frac{\lambda(q'' + q') - 2}{4} T - \frac{\lambda^2}{24} T^3 \right], \end{aligned} \quad (4.9)$$

(ii) the harmonic oscillator with a time-dependent frequency [12], [13]

$$\begin{aligned} \mathcal{K}_v(x'', t''; x', t') &= \lambda_v(2 \sin(\gamma'' - \gamma')) \mid \frac{\sqrt{\dot{\gamma}'' \dot{\gamma}'}}{\sin(\gamma'' - \gamma')} \mid_v^{\frac{1}{2}} \\ \mathcal{X}_v \left[\frac{1}{2} \left(\frac{\dot{s}' x'^2}{s'} - \frac{\dot{s}'' x''^2}{s''} \right) \right] \mathcal{X}_v \left[- \frac{\dot{\gamma}'' x''^2 + \dot{\gamma}' x'^2}{2 \tan(\gamma'' - \gamma')} + \frac{x'' x' \sqrt{\dot{\gamma}'' \dot{\gamma}'}}{\sin(\gamma'' - \gamma')} \right], \end{aligned} \quad (4.10)$$

where $\gamma' = \gamma(t')$, $\gamma'' = \gamma(t'')$ and $s' = s(t')$, $s'' = s(t'')$.

5. Concluding remarks

In this paper we defined p-adic generalization of the Feynman path integral in quantum mechanics. Starting from the definition we calculated it for a particle in a constant external field. Calculation performed simultaneously for real and p-adic cases, and all expressions have the same form with respect to $\mathbb{R}$ and every $\mathbb{Q}_p$.

When classical action $\bar{S}(x'', t''; x', t')$ in the real case is quadratic in x'' and x' there is a general solution [8] which, written in our notation, reads

$$\mathcal{K}_\infty(x'', t''; x', t') = \lambda_\infty \left(-2 \frac{\partial^2 \bar{S}}{\partial x'' \partial x'} \right) \mid \frac{\partial^2 \bar{S}}{\partial x'' \partial x'} \mid_\infty^{\frac{1}{2}} \mathcal{X}_\infty(-\bar{S}(x'', t''; x', t')) .$$

It is natural to expect that the general formula holds

$$\mathcal{K}_v(x'', t''; x', t') = \lambda_v \left(-2 \frac{\partial^2 \bar{S}}{\partial x'' \partial x'} \right) \mid \frac{\partial^2 \bar{S}}{\partial x'' \partial x'} \mid_v^{\frac{1}{2}} \mathcal{X}_v(-\bar{S}(x'', t''; x', t'))$$

whenever $\bar{S}(x'', t''; x', t')$ is quadratic in x'' and x' . Making the Taylor expansion of S around its classical (extremal) value $\bar{S}$ and using the condition (1.3), we can show that $\mathcal{K}_v$ has the above form, but existence of the very prefactor $\lambda_v \left(-2 \frac{\partial^2 \bar{S}}{\partial x'' \partial x'} \right)$ still needs a rigorous proof.

References

- [1] V. S. Vladimirov, I. V. Volovich, E. I. Zelenov, *p-Adic Analysis and Mathematical Physics*, World Scientific, Singapore, 1994.
- [2] L. Brekke, P. G. O. Freund, Physics Reports **233** (1993),1-66.
- [3] A. Khrennikov, *p-Adic Valued Distributions in Mathematical Physics*, Kluwer Acad. Publishers, Dordrecht, 1994.
- [4] W. H. Schikhof, *Ultrametric Calculus: An Introduction to p-Adic Analysis*, Cambridge Univ. Press, Cambridge, 1984.
- [5] V. S. Vladimirov, I. V. Volovich, Commun. Math. Phys. **123** (1989), 659-676.
- [6] B. Dragovich, Int. Journal of Modern Phys. **A 10** (1995), 2349-2365.
- [7] R. P. Feynman, Rev. Mod. Phys. **20** (1948), 367-387; R. P. Feynman, A. Hibbs, *Quantum Mechanics and Path Integrals*, McGraw Hill, New York, 1965.
- [8] C. Morette, Physical Review **81** (1951), 848-852.
- [9] Journal Math. Phys. **36** (5) (1995), a special issue on functional integration.
- [10] E. I. Zelenov, Journal Math. Phys. **32** (1991), 147-152.
- [11] J. J. Halliwell, J. Louko, Physical Review **D 39**(1989), 2206-2215.
- [12] D. C. Khandekar, S. V. Lawande, Journal Math. Phys. **16** (1975), 384-388.
- [13] G. S. Djordjevic, B. Dragovich, *Proc. of Conf. QS96 Minsk*, World Scientific, Singapore, 1996.